

# Computer Aided Molecular Design



**A Teaching License for the classroom use of MOE has been kindly granted by  
the Chemical Computing Group**

1010 Sherbrooke St. West

Suite 910

Montreal, Quebec

H3A 2R7

Canada

**[www.chemcomp.com](http://www.chemcomp.com)**

## **Computer Aided Molecular Design**

### **Introduction**

Our society is faced with challenges that can have a chemical solution. Examples include: bacterial drug resistance, new diseases like AIDS, and agricultural pest control. We also need to develop environmentally benign synthetic methods, for example the use of aqueous solvents and enzymatically catalyzed processes. We are rapidly developing new technologies to help solve these problems, including new receptors such as zeolites and self-assembling systems, and new hosts for guest-host chemistry including crown ethers, cyclodextrins, phase transfer catalysis agents, and supramolecular hosts. Combinatorial synthesis is accelerating the pace of advancement in guest-host chemistry and biochemistry. Each of these areas underscores the need for enhanced uses of molecular mechanics and dynamics, molecular orbital calculations, and chemical information technology.

We are part of a large scale international effort to solve the many problems that confront our society. In this effort we have amassed a truly bewildering array of information; we need to learn how to tame this rich resource. A tremendous volume of work has been done, and a bewildering variety of structural motifs have been discovered in nature. Chemical information technology helps us to appreciate the richness and variety of chemical structural complexity. Computer Aided Molecular Design, CAMD, is a combination of computational chemistry and information technology tools that help us to discover new and useful compounds.

CAMD touches all areas of chemistry. The discovery of new natural products help us explore structures and functionality that we would never guess are important, for example diepoxides and diacetylides. The vastness of the variety of chemical structures also plays an important role in environmental chemistry and analytical chemistry. Characterizing the biological activity and properties of all the known compounds is impossible. We must develop predictive tools for molecular properties in an environmental setting. Quantitative structure activity relationships (QSAR), Quantitative structure property relationships (QSPR), and 3D-database mining play a central role in this effort. Analytical chemists have developed new chemometric techniques that allow the rapid retrieval and prediction of molecular and biological properties. Multi-variate and artificial intelligence techniques are necessary to efficiently use our wealth of information.

This tutorial introduces the use of advanced computational methods in Computer Aided Molecular Design. Computer Aided Molecular Design (CAMD) is a unifying theme that focuses on why we do chemistry and how we decide what to synthesize and study. Chemistry emphasizes the development of predictive tools for understanding structure-function relationships, and the use of CAMD techniques enhances our ability to predict chemical reactivity and design useful compounds.

### **Computer Aided Molecular Design Phases**

The goal of Computer Aided Molecular Design (CAMD) is to find ligands that are predicted to interact strongly with a host. Alternatively, this procedure can be reversed to search for hosts that will interact strongly with a given ligand. CAMD is an outgrowth of rational drug design<sup>1</sup> where the interactions are protein or DNA binding with substrates. It is clear, however, that CAMD is not restricted to drug design. In fact, much of the current development in chemistry, biochemistry, and biology are coalescing and the divisions in our disciplines are evaporating. As a consequence the tools developed for drug design will become critical for many if not most chemists. In particular, molecular recognition, be it through proteins, DNA, supramolecular chemistry, or self-assembling systems, is a unifying research area. As organic and physical

chemists search for guest-host systems with specificity in binding and catalysis<sup>2,3</sup>, the basic concepts of molecular field analysis and receptor mapping will be a unifying tool. Rapid advancements in chemistry will increasingly require an interdisciplinary approach; biochemistry, molecular biology, microbiology, cell biology, developmental biology will be key players along with the traditional areas of chemistry.

The ready availability of chemical structure databases is playing an important role in enhancing the drug discovery approach and CAMD. These same databases find increasing use in environmental, inorganic, and organic chemistry. Supramolecular and natural products chemistry will demand easy access to information and powerful 3D-searching algorithms. Environmental chemists will need to systematize the reactivity of millions of naturally occurring and synthetic substances.

The basic phases of CAMD can be outlined as shown in Table I<sup>1,4</sup>.

Table I. The Phases in Computer Aided Molecular Design<sup>1,4</sup>. The "CPU " column compares the relative computation resources needed for each method.

Phase	Method	CPU	
<b>Determine structure of the ligands or the receptor site:</b>	MO calculations	+	
	molecular mechanics	+	
	molecular dynamics/protein folding	+++	
	homology modeling with database	+++	
↓			
<b>Build a model of the receptor site:</b>	propose pharmacophore	3D-QSAR or receptor mapping	++
	propose steric pocket	map surface with a probe	+
		steric model from map (DOCK)	+
↓			
<b>Search databases for ligands:</b>	2D-substructure	+	
	steric search (docking)	++	
	3D-search with pharmacophore	+++	
↓			
<b>Dock new ligands to receptor site:</b>	molecular mechanics or MO	+	
↓			
<b>Predict binding constants or activity:</b>	1D, 2D, or 3D-QSAR	+	
	free energy perturbation	+++	
	MO transition-state calculations	+++	
↓			
<b>Synthesize ligands:</b>	reactions database	+	

CAMD can be done in two ways: ligand based or receptor based. Receptor based design starts with a known receptor, such as a protein binding site or supramolecular host. Ligand based

design uses a known set of ligands, but an unknown receptor site. Both approaches are actually very similar.

*Receptor based CAMD:* The first phase is to determine the structure of the binding site using standard structural analysis from X-ray diffraction, NMR, or calculations involving molecular orbital or molecular mechanics and dynamics techniques. In the absence of structural information, homology of the unknown receptor sequence with known structures that have been identified through database searches may be a good starting point.

The next phase is to generate a query for database searching. The query is generated by building a simplified model of the receptor site. This model may be based on a pharmacophore, which identifies a few specific interactions that are responsible for the binding (Figure 1.). These interactions include hydrogen bond donors and acceptors, charged groups, and hydrophilic regions such as hydrocarbon side chains, and phenyl groups. The pharmacophore can be generated by visual inspection or by computational techniques. In docking-based searches, the model is based on an analysis of steric interactions over the receptor site. Typically, a solvent accessible surface map is generated and binding pockets are identified on the host surface. More specific interactions can also be specified as in the angiotensin converting enzyme inhibitor pharmacophore<sup>5</sup> in Figure 1.




Figure 1. Pharmacophore for angiotensin converting enzyme (ACE) inhibitors, which describes the spatial arrangement of functional groups necessary for binding to the receptor site of the enzyme<sup>5</sup>. The dotted bonds are to allow single or double bonds and the A stands for any atom.

The next phase is to search databases for ligands that may bind to the chosen receptor. The 3D-pharmacophore is used in conformationally flexible searches for ligands that match the spatial distribution of the receptor. Alternatively, the receptor pocket can be used with auto-docking to find ligands that avoid close-contacts. The 3D-pharmacophore approach and the binding pocket approach are actually very similar, and queries can be fashioned that incorporate aspects of both approaches. Pharmacophores emphasize a few specific and varied types of interactions, while binding pockets emphasize steric interactions over the entire ligand.

The results of the database search may be used directly or modified to produce candidates for further study. These candidates that are inspired by the database search constitute the design element of the procedure. The new ligands or hosts are then assessed for the use at hand. This assessment first involves docking the new molecule and evaluation of the full interaction by molecular orbital calculations or molecular mechanics. Next, calculations are done to predict the binding constant or activity of the compound.

Prediction of the binding constants are usually performed using Gibbs free energy perturbation studies based on either Monte Carlo or molecular dynamics simulations. Activity predictions are usually based on QSAR extrapolation. Increasingly these QSAR predictions are based on the 3D-QSAR that was used to generate the pharmacophore in the search stage.

Finally, the candidates are synthesized and tested in the laboratory. Synthetic chemists increasingly use reaction database searches and artificial intelligence tools to design synthetic procedures.

*Ligand based CAMD:* Ligand based design starts with a group of ligands that have known binding constants or biological activities. The first phase is to determine the structure of the ligands using standard structural analysis from X-ray diffraction, NMR, or calculations involving molecular orbital or molecular mechanics and dynamics techniques.

The next phase is to generate a query for database searching. The query is generated by building a simplified model of the receptor site. This model is based on a pharmacophore, as in receptor based design. The pharmacophore can be generated by visual inspection or by statistical techniques. One popular statistical technique is 3D-QSAR as represented by the CoMFA approach<sup>6</sup>. 3D-QSAR maps the steric, charge, and hydrogen bonding interactions into a 3-D grid for each known ligand. These maps are then compared to find features that the active compounds have in common. The map of common features is then converted into a pharmacophore.

The next phase is to search databases for new ligands that may also bind to the chosen receptor. 2D-substructure searches based on the known ligands can be used, but such searches have not been very successful. Instead, the 3D-pharmacophore is used in conformationally flexible searches for ligands that match the spatial distribution of the known ligands.

The remainder of the phases are identical for ligand and receptor based design.

There are many examples of applications of CAMD. Organic chemists are active users of database searching. Physical chemists calculate the free energy of binding or solvation for substances by perturbation techniques. QSAR/QSPR is a major focus in physical organic chemistry, instrumental analysis, and environmental chemistry. Biochemists focus on receptor modeling and substrate docking. In synthetic chemistry reaction databases play an important role. In inorganic chemistry, 3D-database searches are applied to organometallic chemistry and metal complexes.

## BIBLIOGRAPHY

1. Y. C. Martin, Computer Assisted Rational Drug Design, in *Methods In Enzymology*, D. M. J. Lilley and J. E. Dahlberg, Eds., Academic Press, San Diego, CA, **1991**, pp 587-613.
2. J. Rebek, Jr., Molecular Recognition and Biophysical Organic Chemistry, *Acc. of Chem. Res.* **1990**, *23*(12), 399-404.
3. Organic 'Tectons' Used To Make Networks With Inorganic Properties. *Chem. Eng. News* **1995**, January 2, 21-22.
4. L. M. Balbes, Guide to Rational (Computer-Aided) Drug Design, Research Triangle Institute, Research Triangle Park, NC 27709-2194, from the Ohio Supercomputer Center Computational Chemistry Bulletin board.
5. D. R. Henry, O. F. Güner, Techniques for Searching Databases of Three-Dimensional (3D) Structures with Receptor-Based Queries, Electronic Conference on Trends in Organic Chemistry ECTOC-1, **1995**, Paper 44, <http://www.ch.ic.ac.uk/ectoc/papers/44/.index.html>
6. R. D. Cramer, III, D. E. Patterson, J. D. Bunce, Comparative Molecular Field Analysis (CoMFA). 1. Effect of Shape on Binding of Steroids to Carrier Proteins, *J. Am. Chem. Soc.*, **1988**, *110*(18), 5959-5967.

## QSAR

### Introduction

The key insight of chemistry is the relationship between molecular structure and molecular function. We use the details of molecular structure to predict the properties of molecules. Medicinal chemistry is a particularly rich example of our use of structure-function relationships. There is a tremendous need to be able to quickly design new drugs for curing human disease. The rapid prediction of the activities of compounds for use as drugs and the discovery of new compounds is an important goal. Quantitative Structure Activity Relationships, or QSAR, allow us to predict the properties of compounds and are a quantitative expression of structure-function relationships. QSAR has been responsible for the rapid development of many new drugs.

In QSAR we seek to uncover correlations of biological activity with molecular structure. With Quantitative Structure Property Relationships, QSPR, we extend the same notion to general chemical property prediction and not just biological activity. In either case, the relationship is most often expressed by a linear equation that relates molecular properties,  $x, y, \dots$ , to the desired activity,  $A$ . For compound  $i$ :

$$A_i = m x_i + n y_i + o z_i + b \quad 1$$

Where  $m, n$ , and  $o$  are the linear slopes that express the correlation of the particular molecular property with the activity of the compound, and  $b$  is a constant. If only one molecular property is important, for example molecular volume, then Eqn. 1 reduces to the simple form of a straight line,  $A_i = m x_i + b$ . The slopes and the constant in Eqn. 1 are often calculated using multiple linear regression, MLR, which is analogous with regular linear regression when there is just one independent variable. The molecular properties can be dipole moments, steric energies, molecular volumes, surface areas, free energies of solvation, and a wide variety of others. The molecular properties used in QSAR studies are called descriptors. In Eqn. 1 we show only three descriptors. In a typical QSAR study, scores of descriptors are often used. However, in the final QSAR equation we seek to find the smallest number of descriptors that can adequately model the activity of the compounds in the study. For the general case with  $p$  descriptors,  $x_j$ :

$$A_i = \sum_{j=1}^p m_j x_j + b \quad 2$$

gives the more general QSAR equation form.

### Activities

An example of a QSAR study is the isonarcotic activity of esters, alcohols, ketones, and ethers with tadpoles, Table 1. In this study various organic compounds were added to water with swimming tadpoles. The swimming speed of the tadpoles was observed and the amount of the compound that was necessary to slow the tadpoles swimming was determined. A very effective compound has a very low concentration for the production of the desired effect. In QSAR studies we often like to have the more effective compounds have a higher "activity," not a lower. Therefore, it is very common to transform the concentration for a desired effect,  $C$ , to an activity by:

$$A = \log(1/C)$$

The  $\log(1/C)$  value increases with compound efficacy.

### Log P

The most common descriptor used in QSAR studies is  $\log P$ , which is the  $\log_{10}$  of the octanol/water partition coefficient:

$$P = \frac{c(\text{octanol})}{c(\text{water})} \quad 3$$

The octanol/water partition coefficient is measured by placing the compound in a separatory funnel with octanol and water. Octanol and water are immiscible, and the compound under study partitions between the two phases. The concentration of the compound in the two phases and hence the partition coefficient are a measure of the hydrophobic-hydrophilic character of the compound. The more hydrophobic, the larger are  $P$  and  $\log P$ .  $\log P$  is a common descriptor in QSAR studies because drugs must often cross membranes. Cell membranes are composed of phospholipids, which have hydrophobic tails that produce a very hydrophobic environment in the middle of the membrane bilayer. In the absence of active membrane transport, more hydrophobic drugs have an easier time getting through a membrane.

Table 1. Isonarcotic Activity of Esters, Alcohols, Ketones, and Ethers with Tadpoles

Compound	$\log(1/C)$	$\log P$
CH <sub>3</sub> OH	0.30	-1.27
C <sub>2</sub> H <sub>5</sub> OH	0.50	-0.75
CH <sub>3</sub> COCH <sub>3</sub>	0.65	-0.73
(CH <sub>3</sub> ) <sub>2</sub> CHOH	0.90	-0.36
(CH <sub>3</sub> ) <sub>3</sub> COH	0.90	0.07
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> OH	1.00	-0.23
CH <sub>3</sub> COOCH <sub>3</sub>	1.10	-0.38
C <sub>2</sub> H <sub>5</sub> COCH <sub>3</sub>	1.10	-0.27
HCOOC <sub>2</sub> H <sub>5</sub>	1.20	-0.38
C <sub>2</sub> H <sub>5</sub> COC <sub>2</sub> H <sub>5</sub>	1.20	0.59
(CH <sub>3</sub> ) <sub>2</sub> C(C <sub>2</sub> H <sub>5</sub> )OH	1.20	0.59
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>3</sub> OH	1.40	0.29
(CH <sub>3</sub> ) <sub>2</sub> CHCH <sub>2</sub> OH	1.40	0.16
CH <sub>3</sub> COOC <sub>2</sub> H <sub>5</sub>	1.50	0.14
C <sub>2</sub> H <sub>5</sub> COC <sub>2</sub> H <sub>5</sub>	1.50	0.31
CH <sub>3</sub> (CH <sub>2</sub> ) <sub>4</sub> OH	1.60	0.81
CH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> COCH <sub>3</sub>	1.70	0.31
CH <sub>3</sub> COOCH <sub>2</sub> C <sub>2</sub> H <sub>5</sub>	2.00	0.66
C <sub>2</sub> H <sub>5</sub> COOC <sub>2</sub> H <sub>5</sub>	2.00	0.66
(CH <sub>3</sub> ) <sub>2</sub> CHCOOC <sub>2</sub> H <sub>5</sub>	2.20	1.05




Figure 1. Isonarcotic Activity of Esters, Alcohols, Ketones, and Ethers with Tadpoles

When the data in Table 1 is submitted to least squares linear regression, Figure 1, the resulting QSAR equation is:

$$\log(1/C) = 0.731 \log P + 1.22 \quad n = 20 \quad r = 0.881 \quad 4$$

The data for 20 compounds is reasonably correlated with a regression coefficient of 0.881, indicating a moderately good fit. In this study only one descriptor is necessary to build an adequate model of the structure-function relationships, but often many descriptors are needed. Addition of other descriptors would certainly improve the fit for this case also.

#### *Correlation does not imply Causation*

It is always important for studies of this type to underscore the difference between correlation and causation. We shouldn't read too much into QSAR equations. A good QSAR correlation does not mean that the particular descriptor "causes" the efficient action of the drug. For example, the correlation of isonarcotic activity to log P does not necessarily mean that getting the drug across the membrane barrier is the important step for biological activity. Instead, the log P dependence may be caused by more efficient blood transport, more efficient interactions with nerve receptors, or a myriad of other interactions both major and minor that add up to the net effect. The lack of evidence on causation, or in other words the mechanism of action, may be disappointing at first. However, the goal of QSAR is to predict activity, and that goal is often admirably filled. Information on the various mechanisms leading to biological activity must be done through additional, careful experimentation.



## The History of QSAR

The genesis of QSAR is from physical organic chemistry and linear free energy relationships. The first such studies were done by L. P. Hammett. His goal was to uncover the effects of electronic structure on organic reactivity. A short discussion of his work will be instructive as we start to understand the foundations of QSAR. Hammett's first studies were to understand the effect of electron withdrawing and donating groups on the pK<sub>a</sub>'s of substituted benzoic acids, Table 2. Hammett first wanted to develop a descriptor that described inductive substituent effects. He compared the log K<sub>a</sub> for a variety of substituted benzoic acids with the log K<sub>aH</sub> for unsubstituted benzoic acid to define the  $\sigma$  substituent constant.


$$\sigma = \log K_a - \log K_{aH} \quad 5$$

Table 2 lists the Hammett constants for meta and para substituents. He then postulated that other properties, other than acidity, would be likewise effected by the same substituent effects, and that these other properties would follow the relationship

$$\log (\text{property}) = \rho \sigma + \text{cst} \quad 6$$

In other words, the expected function was a linear relationship with the  $\sigma$  inductive constant. Hammett showed that a wide variety of organic thermodynamic and kinetic properties followed Eqn. 6.

Table 2. Hammett substituent constants for inductive effects.


	Group	$\sigma_p$	$\sigma_m$
	-NH <sub>2</sub>	-0.57	-0.09
	-OH	-0.38	0.13
	-OCH <sub>3</sub>	-0.28	0.10
	-CH <sub>3</sub>	-0.14	-0.06
	-H	0	0
	-F	0.15	0.34
	-Cl	0.24	0.37
	-COOH	0.44	0.35
	-CN	0.70	0.62
	-NO <sub>2</sub>	0.81	0.71

Why did Hammett choose to work with log functions? He took his inspiration from the relationship between Gibbs Free Energy and the equilibrium constant for a reaction,

$$\Delta_r G = - RT \ln K_{eq} \quad 7$$

In other words, the energetics of reactions is related to concentration measurements by logarithmic relationships. Eqn. 6 is just one example of many such relationships used by Hammett and many other organic chemists. All such equations are called linear free energy relationships, or LFERs. In this context, QSAR is just an extension of LFERs into the world of biological structure-function relationships. Corwin Hansch at Pomona College is responsible for the early development of QSAR, and has compiled an extensive database of thousands of QSAR equations that are extracted from the literature (<http://clogp.pomona.edu/>). Hammett  $\sigma$  constants are commonly used as descriptors in QSAR.

An example of using Hammett  $\sigma$  constants in QSAR is work on sulfa drugs. The activity of different sulfa drugs was tested on *E. coli*. Linear regression yielded the QSAR equation



Therefore, even though Hammett's  $\sigma$  constants were derived from acidity data of benzoic acids, the same inductive effects correlate well with the activity of a common class of antibiotics. The important point of Eqn. 8 is that we now have a hint of how to design a better drug: choose a substituent with a larger  $\sigma$  or use multiple substitutions.

### Descriptors

A common issue in chemistry is how to describe molecules and their properties. Quantitative measures are necessary for QSAR studies. Molecular descriptors fall into three general categories, steric, hydrophobic, and electronic. Molecular volumes, surface areas, and bond connectivity indices are steric in nature. Molecular volume and surface area are calculated by placing a sphere on each atom with the radius given by the Van der Waals radius of the atom. The steric energy from molecular mechanics calculations is quite useful. Another useful steric descriptor is the number of rotatable bonds. As the number of rotatable bonds increases, the molecule becomes more flexible and more adaptable for efficient interaction with hosts.

Electronic descriptors include dipole moments, polarizability, and electronic energies. These values are available from molecular orbital calculations. The HOMO-LUMO band gap energy is often used for QSAR. Hammett substituent constants are also examples of electronic descriptors. The number of hydrogen bond donors and acceptors and measures of the pi-pi donor-acceptor ability of molecules are also useful.


Log P is an example of a hydrophobic molecular descriptor. Molecular surface areas can also be divided into hydrophobic and hydrophilic parts. For these surface area descriptors, Van der Waal's spheres are again used, but the hydrophilic surface area is the sum over all electronegative surface atoms (N, O, S) and H's in polar bonds (e.g. O-H, N-H, S-H). The hydrophobic surface area is the sum over all non-electronegative atoms and H atoms that are in non-polar bonds ( e.g. C-H bonds). Log P is also the basis for other descriptors.

### $\pi$ : Lipophilic Character

The lipophilic character,  $\pi$ , of a substituent is defined as

$$\pi = \log P - \log P_H \quad 9$$

where  $\log P$  is the value for the substituted benzoic acid and  $\log P_H$  is the value for unsubstituted benzoic acid.  $\pi$  is a commonly used descriptor for hydrophobic or lipophilic effects. The concentration of substituted penicillins curing 50% of mice infected with *S. aureus* was determined and the activity correlated well with  $\pi$ :




$$\log 1/C = -0.46 \pi + 4.85 \quad n=20 \quad r=0.91 \quad 10$$

### Free Energies of Desolvation


Solvation plays a very important role in molecular interactions.  $\log P$  is directly related to solvation.  $P$  is the equilibrium constant for the process:



and  $\Delta_p G$  is the corresponding Gibbs Free energy change for the process. The partitioning process can be broken into two separate processes, the desolvation from water



and the desolvation from octanol:



The Free Energy of desolvation in water,  $\Delta_{\text{desol}} G(\text{aq})$ , is often written as  $F_{H2O}$  or  $E_{\text{sol}}$  and the Free Energy of desolvation in octanol,  $\Delta_{\text{desol}} G(\text{octanol})$ , is often written as  $F_{\text{oct}}$ . Eqn. 11 for the full partitioning is equivalent to Eqn. 12 – Eqn 13, and the energy effects are

$$\begin{aligned} \Delta_p G &= \Delta_{\text{desol}} G^\circ(\text{aq}) - \Delta_{\text{desol}} G^\circ(\text{octanol}) \text{ or} \\ \Delta_p G &= F_{H2O} - F_{\text{oct}} \end{aligned} \quad 14$$

$\log P$  can then be calculated using Eqn. 11:

$$\log P = - (1/2.303RT) F_{H2O} - F_{\text{oct}} \quad 15$$

Note that  $1/2.303RT = -0.735 \text{ kcal}^{-1} \text{ mol}$ . Therefore, if experimental values for  $\log P$  are not known,  $\log P$  can be estimated from free energies.

Free energies of desolvation have been studied extensively. A group additive properties approach has been taken to estimating these free energies. Group additive property, GAP, approaches are common in estimating thermodynamic variables. For GAP calculations we assume simply that a molecule is the sum of its parts. Table 2 lists several GAP parameters for Fh2o and Foct estimation. These contributions are summed to get the Fh2o and Foct for the entire molecule.

Table 2. Group Additive Property parameters for Free Energies of Desolvation.

Group	Fh2o (kcal/mol)	Foct (kcal/mol)
CH <sub>3</sub> (methyl-aliphatic)	0.800	-0.160
CH <sub>2</sub> (methylene)	0.200	-0.520
CH (methane)	-0.240	-0.560
O (ether-aliphatic)	-3.970	-1.810
O (hydroxyl-aliphatic)	-5.820	-3.790
N (amide)	-2.930	0.020
H (amide)	-3.030	-3.490
Cl (aliphatic)	-0.940	-1.020

For example, the Gibbs Free Energy of desolvation for dimethylether from aqueous solution is estimated to be 0.800+0.800+-3.970 kcal/mol.

GAP estimation makes Free Energies of desolvation very easy to calculate for use in estimating log P. Fh2o and Foct are also good molecular descriptors in themselves, and are commonly used directly in QSAR studies.

#### *Atomic Polarizability, APOL*

The polarizability of a molecule is a measure of the ease with which the electron cloud of the molecule can be distorted by an applied electric field. The attractive part of the Van der Waals interaction is a good measure of the polarizability. Highly polarizable molecules can be expected to have strong attractions with other molecules. The polarizability of a molecule can also enhance aqueous solubility. Remember that the Lennard-Jones form of the Van der Waal's interaction is given as (please review the Molecular Mechanics Tutorial):

$$E_{vdw,ij} = -\frac{A}{r_{ij}^6} + \frac{B}{r_{ij}^{12}}$$

In QSAR studies the molecular polarizability is often approximated by the sum of the Lennard-Jones A coefficients for every atom in the molecule, APOL.

#### *Molar Refractivity*

Molar refractivity, MR, is one of the oldest and most successful descriptors for QSAR studies. MR often shows a strong correlation with ligand binding. Both log P and MR increase with alkyl chain length, so log P and MR show a strong correlation. However, polar functional groups increase MR, but decrease log P. Therefore, some authors suggest that MR is a measure of non-lipophilic interactions, while log P is a measure of lipophilic interactions. MR is defined by the Lorentz-Lorentz equation:

$$MR = \frac{(n^2 - 1)}{(n^2 + 2)} \left( \frac{MW}{d} \right)$$

Where  $n$  is the index of refraction,  $MW$  is the molecular weight, and  $d$  is the density.  $MR$  has a strong correlation with the molecular polarizability. When experimental values are not available,  $MR$  is also often approximated from group-additive constants, like many other descriptors.

### Summary

Physical chemistry has often been called “the elegant science of drawing straight lines.” This tongue-in-cheek saying is never truer as in QSAR. In one sense there is nothing magical about searching for log-linear relationships in biological activity. The QSAR results don’t really explain anything. QSAR equations just point to correlations. On the other hand, QSAR is a very important and routine method for many areas of chemistry. QSAR is best appreciated as a guide for our chemical intuition. QSAR models guide us to what to synthesize next in our search for more effective solutions to our problems.

## Design and QSAR

One goal of QSAR is to decide what molecules to synthesize next. It is sometimes difficult to decide how to make changes to lead compounds based on the QSAR coefficients and descriptors. Some cases are easy. If molecular weight has a strong positive correlation with activity, simply increasing the molar mass by adding some heavy substituents like Br or *t*-butyl may work well. If surface area is an important component then adding a benzene ring might be useful. If log P has a strong positive correlation, then adding a “greasy” side chain like an *n*-butyl or cyclohexane ring would increase lipid solubility.

One strategy of medicinal chemists is the use of the series: methyl, ethyl, propyl, isopropyl, butyl, *t*-butyl for substitutions at various spots in the molecule. For the series methyl, ethyl, propyl, and *t*-butyl, molecular weight, surface area, molar refractivity, log P, and  $\pi$  increase, Table 1. The volume varies as: methyl, ethyl, propyl ~ iso-propyl, *n*-butyl, *t*-butyl. Halogens are also commonly substituted for hydrogen. Fluorine has similar hydrophobic character to hydrogen, but a larger surface area.

Table 1. Substituent Parameters for Common Substituents.<sup>1</sup>

Substituent	Volume (SA)	MR	$\pi$	Rot Bonds
-H	1.48	0.10	0 (reference)	0
-CH <sub>3</sub>	18.78	0.57	0.56	0
-CH <sub>2</sub> CH <sub>3</sub>	35.35	1.03	1.02	1
-CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	51.99	1.5	1.55	2
-CH(CH <sub>3</sub> ) <sub>2</sub>	51.33	1.5	1.53	1
-CH <sub>2</sub> CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	68.63	1.96	2.13	3
-C(CH <sub>3</sub> ) <sub>3</sub>	86.99	1.96	1.98	1
-C <sub>6</sub> H <sub>5</sub>	72.20	2.54	1.96	1
-F	7.05	0.10	0.14	0
-Cl	15.85	0.60	0.71	0

Both log P and MR increase with alkyl chain length, so log P,  $\pi$ , and MR show a strong correlation. However, polar functional groups increase MR, but decrease log P. MR, however, is insensitive to chain branching. The number of rotatable bonds increases ligand flexibility to adapt to a particular binding pocket. Linear alkyl chains have more rotatable bonds than branched alkyl chains. If you want to design in some shape flexibility, use straight chain substituents.

The comparisons in Figure 1 may also be helpful for getting a better feel for log P. LogP increases with alkyl chain length and decreases with chain branching. In other words, branched hydrocarbon chains are more soluble in water than linear chains. Molecules with charged functional groups, like quaternary amines and carboxylates are the most soluble and therefore have very negative logP. Alcohols are more soluble in water than corresponding amines, that is, alcohols have lower log P.






Figure 1. Correlation of log P with structure for alcohols and amines. Calculations by CLOGP program, <http://www.daylight.com/release/index.html>.

The concept of biological isosteres is also useful when designing ligands. Functional groups that have similar properties are called isosteres. The similarity can be based on size, electronic properties, or chemical function. For example, isosteres for the carboxyl group are given in Table 2. For the carboxyl isosteres the similarity is based on chemical function, specifically, acid-base characteristics. Size, molar refractivity and log P ( $\pi$ ) have a large variation. The tetrazole group, which is listed third in the table, is one of the most widely used isosteres.

Table 2 Bioisosteres for the carboxyl group<sup>1</sup> with typical pKa values for aryl compounds.<sup>2</sup>

Substituent	pKa	$\pi$	MR
-COOH	4.2	-0.32	6.9
-COO <sup>-</sup>		-4.36	6.0
 (tetrazols)	4.5	-0.48	15.6
		-3.55	14.6
-SO <sub>3</sub> <sup>-</sup> (sulfonates)		-4.76	~10
-SO <sub>2</sub> NH <sub>2</sub> (sulfonamides)	10	-1.82	12.3
-P(O)(OH) <sub>2</sub> (phosphonic acids)	1.42	-1.59	12.6





Sulfonates are completely ionized at physiological pH, so they are a good substitute for carboxyl if the anion form of the drug is the active form. Sulfonamides are very weak acids; therefore they are good isosteres for carboxyl groups only if the active form is protonated. The order of hydrophobicity for the anions is  $-\text{SO}_3^- < -\text{COO}^- < \text{tetrazoyl}$ . If the anionic form of the drug is the bound form and QSAR shows hydrophobic character is better, then tetrazoyl substitution for carboxyl is a good suggestion.

Bioisosteres for the carbonyl group include amides, sulfoxides, sulfones, and sulfonamides. Isosteres for the hydroxyl group include many of the same functional groups, Table 3. The activity of a particular functional group depends on the target protein. For some binding sites

sulfonamides may be good isosteres for carboxyls. For other binding sites sulfonamides might be good isosteres for hydroxyls. In other words, not all isosteric functional groups are interchangeable for every problem. The actual activity of a given ligand always needs to be measured in the laboratory.

When looking at a set of known active compounds for a given pharmaceutical target, the student might wonder how anyone came up with the substitutions that are used. The isosteres we have discussed, however, should be very helpful when looking at a set of active compounds. Using these ideas, you can follow the logic of the medicinal chemists who designed the drugs that are in use today. Example analgesics are shown in Figure 2.

Table 3. Possible bioisosteres for OH.<sup>1</sup>

Substituent	$\sigma_p$	$\pi$	MR
-OH	-0.37	-0.67	2.9
 (amides)	0.00	-0.97	14.9
 (sulfonamides)	0.03	-1.18	18.2
-CH <sub>2</sub> OH	0.00	-1.03	0.72
 (ureas)	-0.24	-1.30	1.37
 (aminonitrile)	0.06	-0.26	1.01




Figure 2. Analgesics showing ethyl and *t*-butyl substitution and a carbonyl or carboxyl isostere.

## References

1. Hansch, C, Leo, A., Exploring QSAR, Fundamentals and Applications in Chemistry and Biology, American Chemical Society, Washington, DC, 1995. Chapt. 13.
2. Franz, R. G., "Comparison of pKa and Log P Values of Some Carboxylic and Phosphonic Acids: Synthesis and Measurement," *AAPS PharmSci.*, **2001**, 3(2), article 10.



### QSAR of Antitumor Activity in Aziridinyl Benzoquinones

Quinones are a general class of compounds with the structure in Figure 1a. These compounds are thought to undergo reduction in cells to form O-quinone methides that are strong alkylating agents<sup>1</sup>. Alkylating agents attach alkyl groups to the bases in DNA. A series of 2,5-bis(1-aziridinyl)-p-benzoquinones have been tested against lymphoid leukemia in mice<sup>2</sup>. The data in our study table are for chronic treatment with daily injection for 12 days and are given as the  $\log(1/\text{MED})$ , where MED is the minimum effective dose that gives a 40% increase in life span. The general formula for the aziridinyl benzoquinones is shown in Figure 1b.




Figure 1a. Benzquinone




Figure 1b. 2,5-bis(1-aziridinyl)-p-benzoquinones

In this exercise we wish to understand the effect of different R groups on the anti-cancer activity of this group of compounds.

### Procedure

1. Double click on the MOE icon on the desktop or program bar. In the MOE window pull down the File menu and choose "Open".

- On Windows systems type in "c:\Documents and Settings\All Users\Documents\moefiles" and press Enter to switch into the proper subdirectory.
- On OSX systems navigate to "Documents/moefiles".

Then click on CWD. This button set sets the current directory as the default directory. Any further access with the file librarian will default to this subdirectory for opening or saving files.

2. Double click on the benzo directory entry. Scroll down and click on the "benzo.mdb" entry. The ".mdb" extension is used for databases. Click on the "OK" button. All 29 of the molecules in this study will be loaded. The experimental activities are already input for you in column 4. The Free Energy of desolvation from water and octanol, Fh2o and Foct, respectively, are also entered for you (See the QSAR Introduction). To make sure that any changes you make in the data set aren't saved under the original file name, we need to save this data set under a different name. Pull down the File menu in the Database Viewer, and choose "Save As..." In the file dialog type in a new name with the following format "newname.mdb" and click "OK".

3. Remember that you can drag downward in one of the molecule cells to change from the molecular name text to a structure. The first entry has  $R_1 = \text{CH}_3$  and  $R_2 = \text{CH}(\text{OCH}_3)\text{CH}_2\text{OH}$  and is the most active compound.

4. To make the structures easier to see, transfer several of the molecules (one at a time) to the MOE window by doing the following. Right click on the molecule in the Database Viewer and select Send to MOE. In the next dialog box select the Clear system before loading data option and click on OK. The purpose of this step is to just visualize the relationships among the various structures; no calculations are necessary. The mouse buttons are mapped as:

PC Mouse buttons: Middle: Reorient molecule—xyz rotation  
Middle and drag in periphery of viewing area: rotate around z only  
Shift-Middle: xyz translation  
Ctrl-Middle: zoom in and out

5. These molecules have already been molecular mechanics minimized. However, we need to make sure that energetic calculations are done with the Merck Force Field, or MMFF94x. Check the Potential button at the lower left. If MMFF94x is not already selected, click the button and choose MMFF94x. We also would like to do solvation energy calculations. Make sure the “Solvation:” option is set to Born. Click on “Fix charges” if the button is active. Click on Apply and Close.

6. We will first try a QSAR using the classic descriptors. In the Database Viewer pull down the Compute menu, slide right on Descriptors, and choose Calculate. In the list of descriptors highlight apol, ASA, b\_rotN, density, dipole, E (potential energy), E\_sol, logP(o/w), mr, and vdw\_vol. Note the definition of each term in the list. Click OK.

7. The first thing to check is the correlation among the different descriptors. We should not keep descriptors that show a strong correlation. Such strong correlations can distort the QSAR equations. Pull down the Compute menu, slide right on Analysis and choose Correlation Matrix. The numbers in the matrix show the correlation coefficients (r). The diagonal elements are 100% since each variable is 100% correlated with itself. The first column gives the correlation of the descriptor with the activity. This type of correlation is good, and gives a strong indication of the usefulness of the descriptor. Clicking one of the coefficients generates a correlation plot for that pair of variables. For example, clicking on a coefficient in the first column produces the scatter plot for the activity versus the chosen descriptor. In other words, the correlation plot is the result of the single variable QSAR for that descriptor. High correlations in the other columns are bad. To see the between-descriptor correlation plot, click on the coefficient for the corresponding pair of descriptors. Look for any correlations (other than the first column) that are greater than about  $\pm 90\%$ . One of each pair of such variables should be deleted unless you have a particularly sound reason to suspect that both variables will be chemically useful.

In this particular exercise, you should find that ASA, mr, apol, b\_rotN, and vdw\_vol are all strongly correlated. Of the four, mr has a large correlation with the activity (note the first column of correlation coefficients). Also mr is one of the most common descriptors after log P for QSAR. Therefore, we will delete ASA, apol, and vdw\_vol and keep mr; see the next step for instructions. Click Close for the Correlation Matrix window.

8. In the Database Viewer, click right on the column heading for ASA and choose Delete. Click on OK in the warning box. Repeat this process with apol, b\_rotN, and vdw\_vol. You should now have columns for eight descriptors.

9. We are now ready to do some curve fitting. Pull down the Compute menu, slide right on Model, and choose QSAR. The QuaSAR-Model dialog window should open. In the list of descriptors highlight all the entries except the activity or act entry. And then click on Fit. The

regression coefficient ( $R^2$ ) is listed in the status line. The QSAR fit equation coefficients should be displayed in the second column. The size of these coefficients does not necessarily indicate the importance of the descriptor since the numerical range of the different descriptor values can be very different. In other words, descriptors with large ranges have smaller coefficients (slopes) for comparable importance in the fit equation. To account for the range of the descriptor, the third column lists the coefficient divided by the standard deviation of the descriptor values.

10. To get the full details on the fit click on the Report button. Note particularly the correlation coefficient. In this example the result is about 0.822, which is a reasonably good fit for this set of descriptors (however, we are using too many descriptors at this point). The full regression equation is listed as:

$$\text{activity} = 5.953 - 0.048 * \text{Fh2o} - 0.085 * \text{Foct} + 0.001 * \text{E} - 0.087 * \text{E\_sol} - 0.215 * \text{mr} + 0.252 * \text{dipole} - 0.032 * \text{density} - 0.251 * \log\text{P(o/w)}.$$

At the bottom of the Results listing you will see the section:

```

RELATIVE IMPORTANCE OF DESCRIPTORS
0.851132 Fh2o
1.000000 Foct
0.002385 density
0.145518 dipole
0.011524 E
0.650641 logP(o/w)
0.534621 E_sol
0.481458 mr

```

The numbers in this section are calculated from the normalized coefficients and show which variables are responsible for the largest portion of the variability in the data set. In this example, Fh2o, Foct, logP(o/w), E\_sol, mr, and the dipole moment are the most important variables. E is the molecular mechanics steric energy, which includes aqueous solvation Gibbs energy. Fh2o and E\_sol are expected to give comparable results, since both are just different ways of approximating the same thing and they have a descriptor-descriptor correlation coefficient of 0.96. Since we have 29 data points we can at best use 4 or at most 5 descriptors in our study (remember that you need at least 5 observations for each new variable). The next step will be to redo the fit with only Fh2o, logP(o/w), and the dipole moment. You can try using E\_sol and or mr in a separate study.

11. Click the go-away box (or pull down the File menu and choose Quit) on the Results window and click on Yes in the subsequent warning. In the QuaSAR-Model dialog highlight the Fh2o, logP(o/w), and the dipole entries. Use steps 9 and 10 above to redo the fit. Note the new correlation coefficient and the RELATIVE IMPORTANCE OF DESCRIPTORS. Note that the relative importance values can change a lot from fit to fit. The correlation coefficient should be a bit smaller, but not significantly smaller. If the correlation coefficient decreases significantly as you leave out descriptors then you used the wrong descriptors in your fit. Click the go-away box (or pull down the File menu and choose Quit) in the Results window.

12. We need to get a visual picture of how well our QSAR fit does in reproducing the activity of the data set. Scatter plots work well for this purpose. Back in the QuaSAR-Model dialog click on the Validate button. Accept the default calculations. Three new columns will be created in the Database Viewer. The predicted activity from the current QSAR equation will be calculated

along with the residual from the experimental value. The Z-score is the number of standard deviations from the mean for the residual for each data point, which is very useful when looking for outliers. Now we can do our plot. Pull down the Compute menu, slide right on Analysis, and choose Correlation plot. Note the instructions just below the menu bar: next click on the heading for the activity column to select the x-axis and then click on the heading for the \$PRED column. The vertical axis displays the predicted activity from the regression equation. If the regression is perfect, the predicted and experimental activities should be identical and the plot will be a straight line with little scatter and a slope of one. If the regression equation is not very good the plot will have a lot of scatter and /or the slope will not be close to one.

13. You can use the residuals or Z-scores in the Database Viewer to look for outliers, which are molecules that appear to act very differently from the others in the study and so fall well off of the regression line. Clicking on the row number in the Database Viewer highlights the observation in the correlation plot.

14. We now need to clean up before doing another regression method. Click away the correlation plot. Click on the headings \$PRED, \$RES, and \$Z-SCORE and successively remove these columns from the spreadsheet. Also delete the columns for E, Foct, and density.

15. Another way to judge the importance of the different descriptors in modeling the variance in the data is to generate a Contingency table. Pull down the Compute menu, slide right on Descriptors, and choose Contingency. In the descriptor list, select the descriptors that you are most interested in and click OK. C, V, U, and R are different statistical measures of which variables are most useful. The definitions and guidelines for their use are given at the top of the printout. Use the values of C, V, U, and R to help decide which variables to use in your final QSAR.

16. Return to steps 9-15 to see if you can develop a better QSAR. Choose the equation that gives the best fit and that is the easiest to interpret in terms of your goals for molecular design. The general rule is that you need five times as many observations as variables for a valid regression. In this case we have 29 molecules (observations) so the maximum number of descriptors (variables) in the fit equation would be five. However, beyond three descriptors regression equations begin to look suspicious. In general **the fewer the descriptors the better** as long as an adequate fit results.

17. Clean up as instructed in steps 14. Quit the Database viewer.

### References:

1. S. P. Gupta, Quantitative Structure-Activity Relationship Studies on Anticancer Drugs, *Chem. Rev.*, **1994**, *94*, 1507-1551.

2. M. Yoshimoto, H. Miyazawa, H. Nakao, Shinkai, M. Arakawa, *J. Med. Chem.*, **1979**, *22*, 491.

## Acetylcholine Esterase QSAR

Cholinergic nerves use acetylcholine as a neurotransmitter. Acetylcholine acts between neurons and in striated muscle tissue at the neuromuscular junction. When acetylcholine binds to the receptor in a nerve cell an ion channel opens that causes a sudden change in  $K^+$  concentration that depolarizes the cell membrane. Before the nerve can fire again, the acetylcholine must be hydrolyzed into an inactive form. The hydrolysis is done by acetylcholine esterase, which is bound in the membrane of the nerve close to the acetylcholine receptor. The reaction is:




Figure 1. The hydrolysis of acetylcholine is catalyzed by acetylcholine esterase (ACE).

Nicotine and acetylcholine share a set of activities based on interaction with the so-called nicotinic receptors in the synapse of nerve cells. Compounds that share similar activities are called agonists. Nicotinic agonist drugs are used to control blood pressure and neuromuscular function by modulating the activity of ACE. The purpose of this exercise is to determine the important properties of compounds that are nicotinic agonists. With these properties in mind we can then decide what new compounds would be better nicotinic agonists.

Table 1. Relative potencies of nicotinic agonists assayed by contracture of frog muscle.

Number	Agonist	Relative potency
1	(+)-anatoxin	110
2	Pyrido[3,4-b]homotropane	~100
3	Isoarecolone methiodide	50
4	3-hydroxyphenylpropyltrimethylammonium	28
5	4-aminophenylethyltrimethylammonium	15
6	3-hydroxyphenylethyltrimethylammonium	9.8
7	Dihydroisoarecolone methiodide	9.1
8	Arecolone methiodide	8.6
9	Coryneine (i.e. dopamine methiodide)	5.3
10	1-Methyl-4-(trifluoroacetyl)piperazine methiodide	4.6
11	1-Methyl-4-acetylpiperazine methiodide	2.6
12	(±)-Isoarecolol methiodide	2.0
13	1-Methyl-4-carbamyl-1,2,3,6-tetrahydropyridine methiodide	0.77
14	1-Methyl-1,2,3,6-tetrahydropyridine-4-methanol methiodide	0.35
15	Dihydroisoarecolol methiodide	0.25
16	1-Methyl-4-carbamylpiperazine methiodide	0.15
17	1-Methyl-4-carbamylpiperidine methiodide	0.052
18	(±)-Octahydro-2-methyl- <i>trans</i> -(1 <i>H</i> )-isoquinolone methiodide	0.015
19	1-Methyl-4-piperidone oxime methiodide	0.0055




Figure 2. Nicotinic agonists.

In this study we will use quantitative structure activity relationships, QSAR, to determine the correlation between structure and function of a set of compounds, Table 1 and Figure 2, that have been tested for contracture of frog muscle<sup>2</sup>. The better the activity the higher the value of the relative potency, which is measured as a log 1/c type value. Here c is the concentration for a particular level of activity.

An important part of this study is the handling of outliers. Often, a large set of data cannot be adequately treated by linear QSAR methods. When this problem occurs, there are three choices: (1) give up, (2) try using more exotic non-linear approaches, (3) remove unusual compounds from the data set. These unusual compounds that cannot be adequately treated by the QSAR are called outliers. After removing outliers, the resulting QSAR is less universal, but the equations can still be used to understand the structure-function relationships of the remaining compounds.

The trick of treating some compounds as outliers is to not discount them completely. After you finish your QSAR study you should go back to look at the outliers. The outliers may in fact be the key to building better solutions to your problem.

### **Procedure**

1. Double click on the MOE icon on the desktop or program bar. In the MOE window pull down the File menu and choose "Open".

- On Windows systems type in "c:\Documents and Settings\All Users\Documents\moefiles" and press Enter to switch into the proper subdirectory.
- On OSX systems navigate to "Documents/moefiles".

Then click on the CWD button. This button sets the current directory as the default directory. Any further access with the file librarian will default to this subdirectory for opening or saving files.

2. Double click on the "nicact" directory in the file list. Scroll down and click on "nicact.mdb" entry. Click on the "OK" button. All 19 of the molecules in this study will be loaded. To make sure that any changes you make in the data set aren't saved under the original file name, we need to save this data set under a different name. Pull down the File menu in the Database Viewer, and choose "Save As..." In the file dialog type in a new name with the following format "newname.mdb" and click "OK".

3. Remember that you can drag downward in one of the molecule cells to change from the molecular name text to a structure. The first entry, anatoxin A, is the most active compound.

4. To make the structures easier to see, transfer several of the molecules (one at a time) to the MOE window by doing the following. Right click on the molecule in the Database Viewer and select Copy to MOE. In the Copy Database Molecule to MOE window select the "Clear system before loading data" option and click on OK. The purpose of this step is to just visualize the relationships among the various structures; no calculations are necessary. They all have a quaternary ammonium group. Locate this charged group. The mouse buttons are mapped as:

PC Mouse buttons: Middle: Reorient molecule—xyz rotation  
 Middle and drag in periphery of viewing area: rotate around z only  
 Shift-Middle: xyz translation  
 Ctrl-Middle: zoom in and out

5. These molecules have already been molecular mechanics minimized. However, we need to make sure that energetic calculations are done with the Merck Force Field, or MMFF94x. Check

the Potential button at the lower left. If MMFF94x is not already selected, click the button and choose MMFF94x. We also would like to do solvation energy calculations. Make sure the “Solvation:” option is set to Born. Click on “Fix charges” if the button is active. Click on Apply and Close.

6. We will first try a QSAR using the classic descriptors. In the Database Viewer pull down the Compute menu, slide right on Descriptors, and choose Calculate. In the list of descriptors highlight apol, ASA, density, dipole, E (potential energy), E\_sol, logP(o/w), mr, and vdw\_vol. Note the definition of each term in the list. Click OK.

7. The first thing to check is the correlation among the different descriptors. We should not keep descriptors that show a strong correlation. Such strong correlations can distort the QSAR equations. Pull down the Compute menu, slide right on Analysis, and choose Correlation Matrix. The numbers in the matrix show the correlation coefficients. The diagonal elements should be 100% since each variable is 100% correlated with itself. The first column gives the correlation of the descriptor with the activity. This correlation is good, and gives a strong indication of the usefulness of the descriptor. High correlations in the other columns are bad. Look for any correlations (other than the first column) that are greater than about 90%. One of each pair of such variables should be deleted unless you have a particularly sound reason to suspect that both variables will be chemically useful. You should find that ASA, mr, apol, and vdw\_vol are all strongly correlated. Of the four, mr has a large correlation with the activity. Also mr is one of the most common descriptors after log P for QSAR. Therefore, we will delete ASA, apol, and vdw\_vol and keep mr. Click Close in the Correlation Matrix window.

8. In the Database Viewer, click right on the column heading for ASA and choose Delete. Click on OK in the warning box. Repeat this process with apol and vdw\_vol. You should now have columns for six descriptors.

9. Pull down the Compute menu, slide right on Model, and choose QSAR. The QuaSAR-Model dialog window should open. In the list of descriptors highlight all the entries except the act entry. And then click on Fit. The regression coefficient ( $R^2$ ) is listed in the status line. The QSAR fit equation coefficients should be displayed in the second column of the list box. The size of these coefficients does not necessarily indicate the importance of the descriptor since the numerical range of the different descriptor values can be very different. In other words, descriptors with large ranges have smaller coefficients (slopes) for comparable importance in the fit equation. To account for the range of the descriptor, the third column lists the coefficient divided by the standard deviation of the descriptor values. Which descriptor has the largest normalized coefficient, and thus explains much of the variability in the data set?

10. To get the full details on the fit click on the Report button. Note particularly the correlation coefficient. In this example the result is about 0.62, which is a poor fit for this set of descriptors. The final fit equation is listed as:

$$\text{act} = -367.98 + 10.22 \text{ density} - 62.57 \text{ dipole} + 0.157 \text{ E} - 5.585 \text{ E}_{\text{sol}} + 11.73 \text{ logP(o/w)} + 31.74 \text{ mr}$$

At the bottom of the Results listing you will see the section:

```
RELATIVE IMPORTANCE OF DESCRIPTORS
0.013005 density
1.000000 dipole
0.201868 E
```



0.738111 E\_sol  
 0.315618 logP(o/w)  
 0.439694 mr

The numbers in this section are calculated from the normalized coefficients and show which variables are responsible for the largest portion of the variability in the data set. In this example, dipole, E\_sol, mr, and logP(o/w) are the most important variables. Since we have 19 data points we can at best use 3 or at most 4 descriptors in our study (remember that you need at least 5 observations for each new variable). In this data set mr and logP(o/w) are fairly highly correlated (0.87), so we should only keep one of them. Since logP(o/w) is the most commonly used descriptor, we will continue with only logP(o/w). Of course in a careful study, a second analysis would also be done with just mr and the better of the two kept. The next step will be to redo the fit with only E\_sol, logP(o/w), and the dipole moment.

11. Click the go-away box (or pull down the File menu and choose Quit) on the Results menu and click on Yes in the subsequent warning. In the QuaSAR-Model dialog highlight the E\_sol, logP(o/w), and the dipole entries. Use steps 9 and 10 above to redo the fit. Note the new correlation coefficient and the RELATIVE IMPORTANCE OF DESCRIPTORS. The correlation coefficient should be a bit smaller, but not significantly smaller. If the correlation coefficient decreases significantly as you leave out descriptors then you used the wrong descriptors in your fit. Click the go-away box (or pull down the File menu and choose Quit) in the Results window.

12. We need to get a visual picture of how well our QSAR fit does in reproducing the activity of the data set. Scatter plots work well for this purpose. Back in the QuaSAR-Model dialog click on the Validate button. Accept the default calculations. Three new columns will be created in the Database Viewer. The predicted activity from the current QSAR equation will be calculated along with the residual from the experimental value. The Z-score is the number of standard deviations from the mean for the residual for each data point, which is very useful when looking for outliers. Now we can do our plot. Pull down the Compute menu, slide right on Analysis, and choose Correlation plot. Note the instructions just below the menu bar: click on the heading for the act column to select the x-axis and then click on the heading for the \$PRED column. In the correlation plot, which variable is plotted on the horizontal axis and which along the vertical axis? A perfect fit would result in the data points lying on a diagonal line with a slope of 1. The correlation plot shows that the most active compound is significantly under-estimated by the QSAR regression equation, and some of the less active molecules are also not well fit.

13. You can use the residuals or Z-scores to look for outliers, which are molecules that appear to act very differently from the others in the study and so fall well off of the regression line. The fourth molecule is better represented by the fit than the first three. Anatoxin-A has the worst fit. Clicking on the row number in the Database Viewer highlights the observation in the correlation plot.

14. We now need to clean up before doing another regression. Click away the correlation plot. Click on the headings \$PRED, \$RES, and \$Z-SCORE and successively remove these columns from the spreadsheet. Also delete the columns for E and density.

15. Since our first attempt was so bad, we need now to turn to some non-traditional descriptors to see if we can get a better fit. Return to the Compute|Descriptors|Calculate dialog and add the following descriptors: b\_rotN, vdW-vol, PEOE-VSA\_PPOS, PEOE-VSA\_HYD, VSA\_base, VSA\_hyd. The vdW and VSA descriptors add up the Van der Waals volumes of the atoms in

different ways. The volumes are based on approximations that don't use the 3D geometry, so you don't need to know the correct conformation of the molecule. Instead the volumes use the connection table and estimate the overlap volumes from bond connectivity. The PEOE descriptors are based on charge assignments using the Gasteiger method. The descriptors are summarized as:

Descriptor	Comment
b_rotN	Number of rotatable bonds
vdw-vol	Total van der Waals volume
PEOE-VSA_PPOS	Total positive polar van der Waals surface area for $q_i > 0.2$ .
PEOE-VSA_HYD	Total hydrophobic van der Waals surface area, for $ q_i  \leq 0.2$ .
VSA_base	Sum of VDW surface areas of basic atoms
VSA_hyd	Sum of VDW surface areas of hydrophobic atoms

Similar descriptions for other descriptors are available from the MOE Help Web pages.

16. Return to steps 9-14 to see if you can develop a better QSAR, but this time you decide which are the best descriptors to use. Record the equation that gives the best fit and that is the easiest to interpret in terms of your goals for molecular design. The general rule is that you need five times as many observations as variables for a valid regression. In this case we have 19 molecules (observations) so the maximum number of descriptors (variables) in the fit equation would be three. In general **the fewer the descriptors the better** as long as an adequate fit results. Remember that mr and vdw\_vol are highly correlated, so don't use both. Check for other highly correlated descriptor pairs.

17. Finally, just do the fit with vdw\_vol, PEOE\_VSA\_HYD, and VSA\_base. Record the coefficients. Did your best fit do better or worse?

19. Notice that none of the fits work out really well. What is the problem with our fits? Look at the scatter plots (see Step 12). What does the position of the data points relative to the fit line tell you? Think about this carefully to make sure you understand the problem. For the answer, look at footnote 3.

20. Clean up as instructed in steps 14.

21. Look at the values for the VSA\_base descriptor in the Database Viewer. Note that only the two most active compounds have a value for this variable. The success of the fit so far has depended on using VSA\_base to account for the unusually large activities for anatoxin-a and pyridohomotropene. Usually we don't like to use a descriptor that has zero values for most of the molecules in the study. These two molecules are just different than the rest. In addition the residual for isoarecolone is particularly poor, since it doesn't have an available basic site, but is still quite active. We will delete anatoxin-a and pyridohomotropene from the data set to see if we can develop a QSAR model that does a better job of describing the activity trends for the remainder of the molecules. Click on the row number "1" in the left most column of the spreadsheet and shift-click on row 2. Now pull down the Display menu, slide right on Entry, and choose Hide Selected. The whole rows should be temporarily removed from the table. In addition, hide all the compounds with activities less than 0.5. You should now have molecules 3 – 13 in the table. Shift and shift-click on all the remaining rows. In the QSAR-Model window, check the "Selected Entries Only" option. Redo your fits including dipole, logP(o/w), and E\_sol

in addition to PEOE\_VSA\_HYD and PEOE\_VSA\_PPOS. With only 11 observations only two descriptors can be justified.

23. Does rejecting these molecules as "outliers" produce a better QSAR as we would normally expect? Is rejecting outliers always a bad idea, or does rejecting outliers help in highlighting fit problems with the remaining observations? Clean up as instructed in step 14.

### **Summary:**

A final comment is that it is difficult to generate a good QSAR for this group of compounds. Some information can be gleaned from our QSAR results, but the results are not what you would like. Another approach to the design strategy is to do receptor modeling, which is the subject of the next exercise.

### **Report:**

1. Give the QSAR equation that best describes the activity trends in this group of compounds with all the molecules present. Keep in mind the rules for the maximum number of descriptors, and the desire to keep the number of descriptors as small as possible. Give the QSAR equation that best describes the activity trends for molecules 3 – 13. Remember the rule on the maximum number of descriptors.
2. Other than the quaternary ammonium cation, what other properties enhance the activity of these compounds? Use your QSAR results to help. For example, if the dipole moment contributes in your QSAR equations from question 1, then you can hypothesize that changing the dipole moment would help. Look at the sign of the fit coefficient for the important variables to decide if a bigger or a smaller value increases activity. Use the "RELATIVE IMPORTANCE OF DESCRIPTORS" listing to decide which descriptors will have the biggest effect. Propose a specific molecule that incorporates these changes; draw the 2D-structure.

### **References and Footnotes:**

1. H. P. Rang, M. M. Dale, J. M. Ritter, P. Gardner, *Pharmacology*, Churchill Livingstone, New York, NY, **1995**, Chapt. 6.
2. T. M. Gund, C. E. Spivak, Pharmacophore for Nicotinic Agonists, in *Methods In Enzymology*, D. M. J. Lilley and J. E. Dahlberg, Eds., Academic Press, San Diego, CA, **1991**. pp 677-693.
3. The most obvious problem is that isoarecolone isn't fit well. Rejecting isoarecolone as an outlier will obviously improve the fit regression coefficient, so we won't bother doing that here. Rather we will focus on another problem. The other problem with our fit is that the most active compounds, anatoxin-a and pyrido-homotropane, are so much more active than the other compounds, that they dominate the fit. Deleting the two most active compounds will allow narrower ranges for the plotted activity axes, which will allow the deviations of the less active compounds to be more easily seen.

## **Building a Receptor Model.**

Often, the active site of an enzyme is unknown. However, it is still necessary to guess the important attributes of the active site to design better drugs. One way to guess the properties of the active site is to assume that the properties of the active site are complementary to active lead drugs. For example, if the active leads have a positive center at a particular location, then the enzyme might have a negatively charged amino acid nearby. If the active leads have a hydrogen bond donor in some spot, then the enzyme might have a hydrogen bond acceptor in a nearby location. While the enzyme is presenting complementary functionality to the leads, it must also minimize steric repulsion and maximize favorable Van der Waals interactions. To guess the shape of the active site that allows such favorable interactions, we look at the 3D- volume occupied by the active leads. The enzyme must not have groups that extend into the volume occupied by the drug. The receptor model surface is then taken as the combined Van der Waals surfaces of the active leads.

Sets of active lead compounds will have important similarities and differences. To determine which similarities are important, we use the experimental activity data of the leads when we build the model. For example, if one of the leads in the study has an additional H-bond donor that the others don't have, we can determine if the extra group is important by looking at the activity data. If the unusual compound has a higher activity than similar leads then the extra group must be important. If the unusual lead has an activity lower than similar compounds then the extra group is not important. Weighting all the lead compounds by their activity reinforces the good attributes of the compounds and de-emphasizes the unimportant features.

Before the receptor model can be built, however, the lead molecules must be aligned so that the active functional groups of the molecules are overlapping in space. This step is often the most difficult step in a receptor study. One reason is that many compounds have rotatable bonds, so that we must find the "active" conformer first and then rotate the molecule to align with the other molecules in the study. For guessing the "active" conformer, we can get some help if some of the drugs have ring systems that restrict the number of possible conformations. We then choose a relatively active lead compound that is inflexible and then find conformations of the more flexible leads that mimic the inflexible ones. Another difficulty in alignment is that usually we don't really know what the active functional groups in the pharmacophore are, so each of the above steps involves some guessing. Often it is necessary to build several receptor models based on alternate assumptions for the pharmacophores.

In this exercise we will use our set of ACE inhibitors to build an active site receptor model for ACE using the Flexible Alignment procedure in MOE. The three most active compounds, with file names anatoxin\_a, isoarecolone-methiodide, and 3-hydphenpropMe3 will be used to build the receptor model. Start up MOE. The Molecular Mechanics Tutor instructions will be very useful as you complete the following steps.

### **Procedure:**

#### **Open the Ligands**

1. Pull down the **File** menu and choose "Open..." If the current directory isn't the /moefiles directory, use the pull down menu at the right of the file name dialog box to choose the "c:\Documents and Settings\All Users\Documents\moefiles" on the PC or "Documents/moefiles" directory on OS-X. Click on the CWD (Current Working Directory)

button if this directory isn't already the default directory. Double click on the nicact directory listing to open that directory. Click on anatoxin\_a.moe and then click on "Open."

2. Similarly open the files isoarecolone\_meth.moe and 3-hyphenpropme3.moe so that all three molecules are shown in the MOE window. They will be overlapping.
3. We must next specify the force field that we wish to use. Check the Potential button at the lower left. If MMFF94x is not already selected, click the button and choose MMFF94x. Make sure the "Solvation:" option is set to "gas phase". Click on Fix Charges if the button is active. Click on Apply and Close.

### Align the Ligands

4. Pull down the Compute menu, slide right on Simulations, and choose Flexible Alignment.
5. To speed the alignment process for this simple exercise, we will choose options that simplify the alignment procedure. This is in part possible because anatoxin-a is rather inflexible, so we don't need to spend a lot of time finding low energy conformations for anatoxin-a. The other two molecules just need to fit to anatoxin-a well. Use the settings shown below.

Alignment Mode:  Flexible  Rigid Body  Refine Existing Alignment

Output Database: flexalgn.mdb

Open Database Viewer

Iteration Limit: 15 Failure Limit: 20 Energy Cutoff: 10

Options:  Calculate Forcefield Charges Prior to Search  
 Stochastic Conformational Search

Click on Settings to choose the following options:

Configuration Limit: 100

Alpha: 1 RMSD Tolerance: 0.5

Gradient Test: 0.01 Maximum Steps: 500

Conformational Search:  Randomly Rotate Single Bonds  
 Randomly Rotate Double and Conjugated Bonds  
 Preserve Chirality of Unconstrained Chiral Centers  
 Randomly Invert Unconstrained Chiral Centers

Atom Based Similarity Terms

<input checked="" type="checkbox"/> Volume	1	<input type="checkbox"/> Aromaticity	1
<input type="checkbox"/> H-Bond Donor	3	<input type="checkbox"/> LogP(o/w)	1
<input checked="" type="checkbox"/> H-Bond Acceptor	3	<input type="checkbox"/> Molar Refractivity	1
<input checked="" type="checkbox"/> Hydrophobe	1	<input type="checkbox"/> Exposure	1
<input type="checkbox"/> Hydrophobe - Don/Acc	-1	<input type="checkbox"/> Partial Charge	1
<input type="checkbox"/> Polar Hydrogens	1	<input checked="" type="checkbox"/> Acid/Base	1

Projected Pharmacophore Similarity Terms

<input checked="" type="checkbox"/> H-Bond Don Projection	1	<input type="checkbox"/> Aromatic Center	1
<input checked="" type="checkbox"/> H-Bond Acc Projection	1	<input checked="" type="checkbox"/> CO2-type Centroid	1

In particular, choosing the Iteration Limit at 15 may miss some possibly good alignments. The functional groups on the molecule allow us to skip looking for Hydrogen Bond donors and Aromaticity. Click on Apply or OK.


6. The database viewer should appear and after a short period the Flexible Alignment should produce 1-6 possible alignments. The results will be sorted by strain energy, U. Often the alignment with the smallest strain energy is best. The Column labeled S lists the "objective function." This value evaluates the quality of the alignment, judged on the basis of the overlap of the specific chemical functionality that you chose for the alignment (H-Bond

donors, Acid/Base, and Volume). Ideally the best alignment will have the lowest strain energy and the highest S value. The first listed alignment will automatically be transferred to the MOE window.

7. Rotate the aligned molecules in the MOE window to observe how the chemical functionality has been aligned. Transfer another alignment or two to the MOE window to see if you agree with the rankings from the S values. Transfer an aligned set of molecules to the MOE window by clicking the right mouse button in the molecule cell. You should see a pop up menu; select Send to MOE. Next you see a new dialog box; select the “Clear system before loading data” and then OK. The new alignment should appear in the MOE main window.

### **Build the Receptor Model**


8. Pull down the Compute menu, slide right on Prepare, and choose Partial Charges..., and select Method: Gasteiger (PEOE) charges. Click on OK. Pull down the Compute Menu and choose Surfaces and Maps, slide right on Interaction (VdW). The bottom half of the Surfaces and Maps window lists the active parameters. Choose Color: Electrostatics Solid as shown below:



Click Create. The sliders control the transparency. The left slider is for the front surface and the right slider for the back surface. Play with these settings to see the effect to get an idea of the spatial relationships. The charge distribution shown is that of the ligand. If we look at the surface as the binding site for the ligand in the protein, the charge distribution for the receptor would be opposite to the ligand. For example, the receptor would have functional groups with partial positive charges placed near the red area of the surface to interact with the negative charge on the carbonyls or the hydroxyl oxygen. Change the Color: setting to ActiveLP. Now the receptor would have corresponding regions to the colors shown. If the ligand surface is hydrophobic (green) the receptor pocket would also be hydrophobic. The new color-coding also shows the locations of the lone pairs on the carbonyls or hydroxyl groups on the ligands. The receptor model would have hydrogen bond donors placed to interact with the lone pairs on the ligand. Change the render setting to Line instead of Solid. This setting will also allow you to see the orientation of the ligands better. Note the location of the carbonyl or hydroxyl oxygens. Click on Delete to clear the surface.

9. In the Surfaces and Maps dialog, change the Surface: to Electrostatic Map. This map is calculated using the Poisson Boltzmann equation (which we will discuss in the electrochemistry portion of the lecture, also see the corresponding chapter in the Molecular Mechanics Introduction). The Poisson Boltzmann equation takes careful account of the charge distribution of the molecule and the mirror charges that are induced in the surrounding solvent. The result is a grid of points with the electrostatic and Van der Waals energy evaluated at each

point around the molecule. The resulting contour surfaces are much more accurate than the electrostatic coloring used for Interaction, or Connolly surfaces, generated in the last step. Click on Create.



The resulting contours are predictions for the type of interactions that might stabilize ligand binding: Hyd for hydrophobic interactions, Acc for hydrogen bond acceptors or regions of positive electrostatic potential on the underlying molecule, and Don for hydrogen bond donors or negative electrostatic potential on the underlying molecule. The sliders control the iso-contour level value (in kcal/mol) for each electrostatic region. Use the slider or the text field to enter the display level. The generated surface will be all points in space for which the interaction potential (electrostatic and van der Waals) is equal to the specified value. The surface will dynamically update (in the MOE window) so that Apply need not be pressed. Note that hydrogen atoms and partial charges *are required* for Electrostatic Maps.

The electrostatic map shows that quaternary ammonium compounds produce a very diffuse (unfocused) electrostatic field. The interaction with this region will be nondirectional. Even though the quaternary ammonium group produces a positive electrostatic field, there is no possibility for interaction with hydrogen bond acceptors, so the red region of your map just highlights the area where negatively charged functional groups might interact and the “Acc” indication is really not applicable for this system. The “Don” and “Acc” indications are particularly applicable, however, when mapping the binding pocket of proteins based on the protein structure directly (rather than in the ligand based approach in this exercise).


### **Finishing Up**

1. In the Surfaces and Maps window, click on the line in the list labeled Electrostatic Map. Then click Delete and then click Close.
2. Click the Close button on the right hand side of the MOE Window.


### Determining a Pharmacophore

Several Angiotensin II antagonists are shown below. These drugs are used instead of the earlier angiotensin converting enzyme inhibitors when side effects are a concern.


#### Angiotensin II Antagonists (log P in parentheses)



Candesartan (5.77)




Irbesartan (6.3)




Valsartan (5.44)


Chiral



Abitesartan (5.56)



Milfasartan (5.68)



Zolasartan (6.26)




A pharmacophore is a search query that will be used to search a database for other compounds that might have similar biological activity. One way to build a pharmacophore is to determine the simplest collection of functional groups that appear in common with a group of active compounds. There is no single correct answer for the pharmacophore.

The best way to determine if a pharmacophore is useful is to use it in database searches. If the database search returns only compounds that are known to have the target biological activity then your pharmacophore is too specific. It is only finding compounds that you already know are active. If your pharmacophore produces thousands of database hits, it is too general. Thousands of hits would be impossible to sort through to determine which would be best for further testing. The best pharmacophore strikes a happy medium. The most useful pharmacophores return a large number of active compounds that are already in use. This ensures that you have chosen functional groups that are appropriate for the purpose. But the most useful pharmacophores also return compounds that have not been tested for biological activity for your target and also display some unusual structural features that you might not have thought of otherwise. In other words the best pharmacophore sparks your imagination about possible lead compounds without leading you down blind alleys.

1. Determine a pharmacophore that would fit all six of these compounds that would provide a minimum number of database search hits of other types of compounds. Be explicit about bond and atom types. Do your searches in the CMC3D database. An easy way to build your pharmacophore is to do a database search for the Class of angiotensin inhibitors to find one of the above compounds, for example

like “%angio%”

will work. Then transfer this compound into Isis/Draw and erase the parts of the molecule that are not common to the other active compounds. Focus on the heteroatom functional groups. Also, the tetrazole group is an isostere for a carboxyl/carboxylate functional group. There are only a few tetrazoles in the database, so keeping the tetrazole in the pharmacophore unnecessarily restricts the possible number of database hits. Use a general carboxyl pharmacophore instead of the tetrazole group, as shown below:




The Q atom type stands for any atom except C and H. The A atom type stands for any atom except H. Find a pharmacophore that recovers all six of the active compounds, but no more than about 30 total compounds. Report this pharmacophore.

2. Erase some additional part of your pharmacophore. Choose a part to erase that you feel would allow much additional flexibility, but at the same time won't generate over 400 additional hits. Report this new pharmacophore and also give an example of a database hit that shows some new functionality not present in the original active compounds that you feel might be worth testing.


## Mutagenicity of Chlorofuranone Derivatives

### Introduction<sup>1</sup>

3-chloro-4-(dichloromethyl)-5-hydroxy-2(5H)-furanone (CIF) is a strong acting bacterial mutagen resulting from the chlorine bleaching of wood pulp. The mutagenicity of CIF has been reported to be in the range of 6,300 induced mutations/nanomole ( $10^{-9}$  mole) in *Salmonella typhimurium* (TA100), making it one of the most potent of the TA100 mutagens. The TA100 test is one of the Ames screening methods used to measure the cancer causing potential of compounds.



3-Chloro-4-(dichloromethyl)-5-hydroxy-2(5H)-furanone (CIF)



CIF Analogs

The mutagenic characteristics of this class of molecules can be traced to interactions between the LUMO orbital of CIF and the HOMO orbital (Highest Occupied Molecular Orbital) of the nucleic acid Guanine in the DNA of the salmonella bacteria. The CIF/DNA interaction prevents proper DNA replication, which results in enzyme mutations in subsequent generations of the bacteria.


Numerous analogs of CIF have been isolated and characterized for their mutagenicity.<sup>2-8</sup> In general these differ in the degree of chlorination and ring substitutions. The CIF analogs to be examined typically have  $R_1 = (\text{H}, \text{OH}, \text{OCH}_3, \text{ or } \text{OCH}_2\text{CH}_3)$ ,  $R_2 = (\text{H}, \text{CH}_2\text{Cl}, \text{CHCl}_2, \text{ or } \text{CH}_3)$ , and  $R_3 = (\text{H or Cl})$ . The mutagenicity is usually expressed as,

$$\ln(\text{TA100}) = \ln\left(\frac{\text{Number of induced mutations}}{\text{Nanomole of CIF analog}}\right)$$

Thus, for the CIF parent structure [3-chloro-4-(dichloromethyl)-5-hydroxy-2(5H)-furanone] the activity can be expressed as  $\ln(\text{TA100}) = \ln(6,300)$  or  $\ln(\text{TA}) = 8.748$ . In the following procedure you will determine the mutagenicity of a CIF analog for which mutagenicity data is unavailable by examining the correlation between the CIF analog, LUMO energy, and the mutagenicity expressed as  $\ln(\text{TA100})$ .

The structures are already drawn and minimized using PM3 for you, to save time. Using PM3 is advisable since the molecules under study have chlorines. We will also consider some more classical QSAR descriptors to help complement the LUMO energy. These descriptors, such as Van der Waals surface area (VSA), molecular volume (Vol), dipole moment, and logP are good indicators of the molecular interactions that aid binding and transport of drugs. The LUMO adds information to the QSAR study on molecular reactivity.

### Chlorofuranone Data Set.



### Procedure Outline (The step-by-step instructions will be given below)

1. Load the furanone database.
2. Fill the database spreadsheet with MOPAC based descriptors and a few general descriptors.
3. Do partial least squares, PLS, based QSAR.
4. Choose the best two-descriptor QSAR equation that uses the LUMO energy.
5. Predict activity of the unknown from the best QSAR equation.

### Procedure

You will need to use the QSAR of Antitumor Activity in Aziridinyl Benzoquinones CAMD tutorial to refresh your memory on how to do a QSAR study. These instructions highlight the differences necessary to do this study.

1. Start up MOE. Load the database that is in the furanone directory, called "furanone.mdb". From the database viewer window, save this database under a different name to avoid corrupting the original data.
2. The molecules should already be entered with activities, except for the last molecule, which is the unknown. The unknown is in the table so that the values for the descriptors will be generated.
3. We just want a few descriptors for this study. Pull down the Compute menu and choose Descriptors. In the descriptors list scroll down and click on PM3\_DIPOLE, PM3\_HF, PM3\_HOMO, PM3\_LUMO. Remember that the Heat of Formation, or HF, isn't a heat of formation really at all. HF is just the total electronic energy of the molecule.
4. We also need to add a few standard, non-quantum mechanical descriptors. Click on VSA, vol, PEOE\_VSA\_HYD, and then SlogP. Click the OK button for the QuaSAR-Descriptor dialog box. After a few minutes, the system will complete a MOPAC calculation on each of the molecules in the spreadsheet, automatically. These steps should result in eight new columns in the spreadsheet.
5. Record the results for the descriptors for the unknown molecule. This molecule is in the last row of the table. You will need these values later.
6. Pull down the Compute menu, slide right on Analysis, and choose Correlation Matrix. Observe the correlations and decide which descriptors are likely candidates for good QSAR models following the guidelines in the QSAR of Antitumor Activity in Aziridinyl Benzoquinones Tutorial. Note that "vol" and VSA have a high correlation so that only one of the two should be used in any QSAR regression. You can delete "vol" from the spreadsheet.
7. Pull down the Compute menu and choose QuaSAR-Model. Choose PLS regression and then do the curve fit. Since we only have 15 observations we should stick to two-descriptor models. Make sure that one of the descriptors is the LUMO energy. Record the resulting R<sup>2</sup> values, as listed in the Status line in the QuaSAR-Model window.
8. Choose the three models that have the best R<sup>2</sup> values. View the reports and record the QSAR equations for these three fits. Remember to include the constant term. Observe the scatter plots of the calculated and experimental activities.
9. Choose the QSAR equation that you feel best models the experimental activities. Use this equation to predict the activity of the unknown.
10. Close down MOE and log off of your computer.

### References

1. The Introduction to this exercise is copied from Peploski, James; Clarkson University CM 132 Laboratory Manual, <http://www.clarkson.edu/~acs15/labs/QSAR.pdf>
2. Montorfani, Sergio; Meier, Pierre; Zimmerli, Bernhard; Jaeggli, Mario; "Mutagenic activity and potential MX contamination of some Swiss drinking waters." *Mitt. Geb. Lebensmittelunters. Hyg.*, **1995**, 86(5), 512-25.

3. Poso, Antti; Tuppurainen, Kari; Gynther, Jukka; "Modeling of molecular mutagenicity with comparative molecular field analysis (CoMFA). Structural and electronic properties of MX compounds related to TA100 mutagenicity," *Theochem*, **1994**, *110*(3), 255-60.
4. Tuppurainen, Kari; Lotjonen, Simo; Laatikainen, Reino; Vartiainen, Terttu; "Structural and electronic properties of MX compounds related to TA100 mutagenicity. A semi-empirical molecular orbital QSAR study." *Mutat. Res.* **1992**, *266*(2), 181-8.
5. Tuppurainen, Kari; Lotjonen, Simo; Laatikainen, Reino; Vartiainen, Terttu; Maran, U.; Strandberg, M.; Tamm, T.; "About the mutagenicity of chlorine-substituted furanones and halopropenals. A QSAR study using molecular orbital indices," *Mutat. Res.*, **1991**, *247*(1), 97-102.
6. Ishiguro, Yasuhisa; Santodonato, Joseph; Neal, Michael W.; "Mutagenic potency of chlorofuranones and related compounds in Salmonella," *Environ. Mol. Mutagen.*, **1988**, *11*(2), 225-34.
7. Ishiguro, Yasuhisa; Lalonde, Robert T.; Dence, Carlton W.; Santodonato, Joseph; "Mutagenicity of chlorine-substituted furanones and their inactivation by reaction with nucleophiles," *Environ. Toxicol. Chem.*, **1987**, *6*(12), 935-46.
8. Ishiguro Y; Santodonato J; Neal M W; "Mutagenic potency of chlorofuranones and related compounds in Salmonella," *Environ. Mol. Mutagen*, **1988**, *11*(2), 225-34.