

## Sperner's Lemma


The purpose of this note is to elaborate the proof of Sperner's Lemma.

**Theorem** (Theorem 2.25). Let  $\mathcal{T}$  be a triangulation of an  $n$ -dimensional simplex  $\Delta$ . Suppose that each vertex of  $\mathcal{T}$  has been assigned a number from the set  $\{0, \dots, n\}$ . Then the number of completely labelled  $n$ -simplices of  $\mathcal{T}$  is odd if and only if the number of completely labelled  $(n-1)$ -simplices of the triangulation  $\mathcal{T} \cap \partial\Delta$  is odd.

The first step of Theorem 2.25 is: Suppose that  $D$  is an  $n$ -simplex of  $\mathcal{T}$ . Then the following statements hold:

- (1)  $D$  is completely labelled  $\Leftrightarrow D$  has exactly one completely labelled  $(n-1)$ -dimensional face.
- (2)  $D$  is not completely labelled  $\Leftrightarrow$  either  $D$  has no completely labelled faces or  $D$  has two completely labelled faces.

Pull apart  $\mathcal{T}$  to obtain a collection  $T$  of disjoint  $n$ -simplices. (See the figure.)



From (1) and (2) we obtain:

- (3) The number of completely labelled  $n$ -simplices in  $\mathcal{T}$  is congruent modulo 2 to the number of completely labelled  $(n-1)$ -dimensional faces of  $T$ .

Now it is clear that:

- (4)  $\{ \text{completely labelled } (n-1)\text{-faces of } T \} =$   
 $\{ \text{completely labelled } (n-1)\text{ faces of } T \text{ on } \partial\Delta \} \cup$   
 $\{ \text{completely labelled } (n-1)\text{-faces of } T \text{ not on } \partial\Delta \}$

Notice that the two sets on the right hand side are disjoint.

Consider an  $(n - 1)$ -face  $F$  of  $T$ . If  $F$  appears on the boundary of  $\Delta$  then it is in exactly one  $n$  simplex of  $T$ . If  $F$  is not on  $\partial\Delta$  then  $F$  appears in exactly two  $n$ -simplices of  $T$ . Thus, the set

$$\{\text{completely labelled } (n - 1)\text{-faces of } T \text{ not on } \partial\Delta\}$$

has an even number of elements. That is

- (5) The number of completely labelled  $(n - 1)$ -dimensional faces of  $T$  is congruent modulo 2 to the number of completely labelled  $(n - 1)$ -faces on  $\partial\Delta$ .

Putting observations (3) and (5) together we obtain

- The number of completely labelled  $n$ -simplices in  $\mathcal{T}$  is congruent modulo 2 to the number of completely labelled  $(n - 1)$ -faces on  $\partial\Delta$ .

This is what we were to prove.

